

Introduction to Distance Sampling

Overview of wildlife population assessment methods

Plot sampling

Distance sampling

Basic idea

Types of distance sampling

Wildlife Population Assessment

- How many are there?
- What are their trends?
- Why?
 - Vital rates (survival, fecundity, etc)
- What might happen if...?
 - Scenario planning
 - Risk assessment
 - Decision support

Methods of estimating abundance

- Complete census
- Plot sampling
- Distance sampling
- Mark-recapture
- Removal method

Complete census


Let

- N = population size (abundance)
- A = size of study region = 5000
- D = animal density = N/A

Method: count everything!

- $N = 412$
- $D = 412/5000 = 0.0824$

Rarely possible in practice!



Plot sampling (or strip transect)

- Let

k = number of strips = 5

L = total line length = $50 \times 5 = 250$


w = the strip half-width = 1

a = area of region covered

$$= 2wL = 2 \times 1 \times 250 = 500$$

n = number of animals counted = 36

- From this, how do we estimate abundance?



Intuitive estimator of abundance

I saw 36 animals

I covered $500/5000 = 1/10^{\text{th}}$ of the study region

So, I estimate there are $36/(1/10) = 36 \times 10 = 360$ animals

$$\hat{N} = \frac{n}{a/A} = \frac{nA}{a} = \frac{36 \times 5000}{500} = 360$$

(Hat “^” means an estimate.)

Concept – Plot sampling

Step 1: How many in covered region, N_a ?

Plot sampling: $N_a = n$

Step 2: Given N_a , how many in study region, N

If transects placed at random:


$$\hat{N} = \frac{N_a}{a/A}$$

Overall: $\hat{N} = \frac{n}{a/A} = \frac{nA}{a} = \frac{nA}{2wL}$ ← for strip transects

Distance (line transect) sampling

- An extension of plot sampling where not all animals in the covered region are detected
- Here
 - $w = 2$ (strip can be wider, as don't have to see everything)
 - $a = 1000$
 - $n = 68$ (more animals seen)
- Let
 - P_a = proportion of animals detected within covered region
- Imagine we know (or can estimate)

$$\hat{P}_a = 0.7$$



Intuitive estimator of abundance

- I saw 68 animals
- The estimated proportion seen was 0.7
- I estimate the true number of animals in the strips was $68/0.7 = 97.1$
- I covered $1000/5000 = 1/5^{\text{th}}$ of the study region
- I estimate there are $97.1/(1/5) = 485.7$ animals

$$\hat{N} = \frac{n / \hat{P}_a}{a / A} = \frac{nA}{a\hat{P}_a} = \frac{68 \times 5000}{1000 \times 0.7} = 485.7$$

Concept – Distance sampling

Step 1: How many in covered region, N_a ?

Distance sampling:
$$\hat{N}_a = \frac{n}{\hat{P}_a}$$

Step 2: Given N_a , how many in study region, N

If transects placed at random:


$$\hat{N} = \frac{\hat{N}_a}{a/A}$$

Overall:
$$\hat{N} = \frac{\frac{n}{\hat{P}_a}}{a/A} = \frac{nA}{a\hat{P}_a} = \frac{nA}{2wL\hat{P}_a}$$
 ← for line transects


- How do we estimate P_a ?

How to estimate P_a ?


- Record perpendicular distance, x , from transect line to each observed object



Estimating P_a



Estimating P_a



$$\hat{P}_a = \frac{\text{area under curve}}{\text{area under rectangle}}$$

Area of rectangle = $12 \times 2 = 24$


Area under curve = $0.25 \times (12 + 11.5 + 11 + 10.5 + 9 + 7 + 4 + 3) = 17$

Therefore

$$\hat{P}_a = \frac{17}{24} = 0.7$$


Distance sampling options

- Types of samples
- Types of measurement
- Types of objects detected
- Types of detection methods




Type of sample Line vs. Point

Line transect




Point transect
(Variable circular plot)



Type of distance measurement


1. Radial vs perpendicular

For line transects, can either measure perpendicular distance from line to object



radial distance and angle

$$x = r \sin(\theta)$$



For point transects
measure radial distance from point to object



Type of distance measurement

2. Exact vs Grouped

Exact distance recorded to each object detected



Distances recorded in intervals



Photo: Rich Guenzel

Type of object

1. Individuals vs Clusters

Each object detected is a single individual



Each object detected is a cluster of individuals
- will need to estimate expected cluster size



Photo: Thomas Norris

Type of Object

2. Direct vs Indirect



Objects are animals (or plants) of interest ...



... or something they produce
(an “indirect survey”)



Another example is a cue count

Method of detection

Active vs Passive

Observers actively search for animals and record distances




Photo: Ullas Karanth

Animals generate their own distances (“passive distance sampling”)



Photo: Steve Dawson



Main ideas thus far

Distance sampling is an extension of plot sampling

In plot sampling, we see everything in the covered region

$$\hat{N} = \frac{n}{a/A} = \frac{nA}{a} = \frac{nA}{2wL \cdot 1} \quad \hat{D} = \frac{\hat{N}}{A} = \frac{n}{2wL \cdot 1}$$

strip transects

In distance sampling, we do not see everything, and we estimate the proportion detected, \hat{P}_a

$$\hat{N} = \frac{n/\hat{P}_a}{a/A} = \frac{nA}{a\hat{P}_a} = \frac{nA}{2wL\hat{P}_a} \quad \hat{D} = \frac{\hat{N}}{A} = \frac{n}{2wL\hat{P}_a}$$

line transects

- How do we estimate P_a ?

$$\hat{P}_a = \frac{\text{area under curve}}{\text{area under rectangle}}$$

line transects

